



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

deep-sea fishes of the world, with a table showing their distribution. A full bibliography and a number of other useful tables are also included.

D. S. J.

Jenkins on Labroid Fishes of Hawaii.—In the *Bulletin of the United States Fish Commission* Dr. Oliver Peebles Jenkins, of Stanford University, gives an account of new species of labroid fishes obtained by him and by others in Honolulu in 1889 and later. The chief collection was made by Dr. Jenkins and his assistant, Dr. George C. Price, under the auspices of De Pauw University. Later, both Dr. Jenkins and Dr. Price were called to Stanford University, and the original collection of fishes, by far the largest yet made about the Hawaiian Islands, was supplemented by others, the principal one being made by Dr. Thomas D. Wood, also of Stanford University.

In the single group of Labridæ and Scaridæ twenty-two new species were obtained. These are described and figured in the present paper. These new species are the following:

<i>Macropharyngodon aquilolo.</i>	<i>Iniistius verater.</i>
<i>Halichæres iridescens.</i>	<i>Cheilinus zonurus.</i>
<i>Halichæres lao.</i>	<i>Pseudocheilinus octotania.</i>
<i>Hemicoris remedius.</i>	<i>Anampses evermanni.</i>
<i>Coris lepomis.</i>	<i>Calotomus irradians.</i>
<i>Hemicoris keleipionis.</i>	<i>Scarus brunneus.</i>
<i>Thalassoma pyrrhovinctum.</i>	<i>Scarus gilberti.</i>
<i>Novaculichthys woodi.</i>	<i>Scarus paluca.</i>
<i>Novaculichthys entargyreus.</i>	<i>Scarus ahula.</i>
<i>Hemipteronotus umbulatus.</i>	<i>Scarus miniatus.</i>
<i>Iniistius leucozonus.</i>	<i>Pseudoscarus jordani.</i>

This list indicates the extreme richness of the Hawaiian fish fauna, its isolation and distinctness as compared with the fauna of the East Indies, and the fact that the few collections yet made about Honolulu have barely touched the wealth of the whole.

D. S. J.

Greene on the Caudal Heart of the Hagfish.—In the *American Journal of Physiology* Dr. Charles Wilson Greene gives his studies on the caudal heart in the California hagfish, *Polistotrema stouti*. This structure was first discovered by Retzius in 1890, who accidentally noticed a paired pulsating organ in the tail of the slime eel (*Myxine*). The function of this structure is to drive the blood of the subcutaneous spaces back into the circulatory system.

We are pleased that Dr. Greene calls this curious animal by its